arXiv:1608.02746v1 [math.OA] 9 Aug 2016

HOLOMORPHIC HILBERT BUNDLES AND THE FRAME EXISTENCE PROBLEM

MOHAMMAD B. ASADI AND ZAHRA HASSANPOUR-YAKHDANI

Abstract. We show that if $A = K(l^2) + \mathbb{C}I_{l^2}$, then there exists a Hilbert A-module that possess no frames.

1. Introduction

The classical frame theory has been generalized to the setting of Hilbert C*-modules by Frank and Larson [5]. They concluded from Kasparov's stabilization theorem that every countably generated Hilbert C*-module over a unital C*-algebra has a standard frame. They asked in [5, Problem 8.1], for which C*-algebra A, every Hilbert A-module has a frame? In 2010, Li solved this problem in the commutative unital case, and characterized the unital commutative C*-algebra A that every Hilbert A-module admits a frame as finite dimensional one [6].

In general case, the conjecture is as follow:

"*Every Hilbert C*-module over a C*-algebra A admits a frame if and only if A is a dual algebra.*"

In the commutative case, Li applied the Serre-Swan theorem. This theorem states that there is an one to one correspondence between finitely generated projective modules over a commutative C*-algebra $\mathrm{C}(\Omega)$ and complex vector bundles over Ω [7].

In [4], Elliott and Kawamura shown that the vector space of bounded uniformly continuous holomorphic sections of every holomorphic uniformly continuous Hilbert bundle of dual Hopf type over pure states of a C*-algebra A admits a unique structure of right Hilbert A-module.

In this paper, we give a partially affirmative response to the above conjecture. Indeed, we have applied the Elliott-Kawamura approach and concluded the following result:

"*If H is a separable infinite dimensional Hilbert space and $A = K(H) + \mathbb{C}I_H$, where $K(H)$ is the C*-algebra of compact operators on H, then there exists a Hilbert A-module that possess no frames.*"

2010 *Mathematics Subject Classification.* Primary 46L08; Secondary 42C15, 46L05.

Key words and phrases. Hilbert C*-modules, frames, holomorphic bundles.

2. Holomorphic Hilbert Bundle

Let A be a C*-algebra, $\hat{A}$ the spectrum of A and $P(A)$ be the set of pure states of A. In general, $P(A)$ is not compact, in this case we consider $P_0(A) = P(A) \cup \{0\}$. However, we set $P_0(A) = P(A)$, when $P(A)$ is compact.

For any $\pi : A \longrightarrow B(H_\pi)$ in $\hat{A}$, and each unite vector $h \in H_\pi$, the linear functional f, given by $f(\cdot) = \langle \pi(\cdot)h, h\rangle$, is a pure state. In this case, we use the notations $\pi = [f]$ and $f = (\pi, e)$, where $e = h \otimes h$. Also, the unitary equivalence class of f (as a set) is equal to

$$R_1(H_\pi) := \{e \in B(H_\pi) : \text{e is a rank one projection}\}.$$

$R_1(H_\pi)$ has a natural holomorphic manifold structure that is independent of the chosen representative element in each equivalence class in $P(A)$ [4]. Therefore, we can identify $P(A)$ as the disjoint union of projective spaces, i. e.,

$$P(A) = \bigcup_{\pi \in \hat{A}} \{\pi\} \times R_1(H_\pi).$$

Then $P_0(A)$ has a natural holomorphic manifold structure and it has a natural uniform structure determined by the seminorms arising from evaluation at the elements of A.

In [4], Elliott and Kawamura introduce the concept of uniformly continuous holomorphic Hilbert bundle of dual Hopf type over pure states of a C*-algebra. In fact, we set

$$\mathcal{H} = \{B(H_\pi, K_\pi)e\}_{(\pi \in \hat{A} \cup \{0\}, e \in R_1(H_\pi))},$$

where K_π is a Hilbert space, for all $\pi \in \hat{A}$. If $X(\mathcal{H})$, the vector space of bounded uniformly continuous holomorphic sections of $\mathcal{H}$, exhausting fibres then the pair $(\mathcal{H}, X(\mathcal{H}))$ is a uniformly continuous holomorphic Hilbert bundle of dual Hopf type. In this case, for any $S \in X(\mathcal{H})$ and any $\pi \in \hat{A}$ there exists a operator $S_\pi \in B(H_\pi, K_\pi)$ such that

$$S((\pi, e)) = S_\pi \circ e \qquad\qquad (e \in R_1(H_\pi))$$

As shown in [4], $X(\mathcal{H})$ is a Hilbert A-module. In fact for any $S, T \in X(\mathcal{H})$, the A-valued inner product is defined by S^*T, where

$$S^*(\pi, e) = e \circ S_\pi^* \in eB(K_\pi, H_\pi), \qquad \text{for all } (\pi, e) \in P_0(A).$$

Since $(S_\pi^* \circ T_\pi)_{\pi \in \hat{A}} \in \prod_{\pi \in \hat{A}}(B(H_\pi))$ is uniformly continuous, we can consider S^*T belongs to A, by a Brown's result [2].

3. Frame Existence Problem

Theorem 3.1. *Suppose that A is a C^*-algebra, $f_0 \in P(A)$, $\pi_0 = [f_0]$, H_{π_0} is a separable Hilbert space and W is a countable subset of $P(A)$ such that $f_0 \in \overline{W} \setminus W$. If there exists a uniformly continuous holomorphic Hilbert bundle of dual Hopf type $\mathcal{H} = (B(H_\pi, K_\pi)e_\pi)_{(\pi,e_\pi)\in P_0(A)}$ such that for any $\pi \in [W]$, K_π is separable and K_{π_0} is nonseparable, then the Hilbert A-module $X(\mathcal{H})$ possess no frames.*

Proof. Assume that $\{S_j\}_{j\in J}$ is a frame for $X(\mathcal{H})$. Hence, there exist positive numbers C, D such that for any section $S \in X(\mathcal{H})$ the following inequality holds

$$CS^*S \leq \Sigma_{j\in J} S^*S_jS_j^*S \leq DS^*S.$$

Hence, for every $\pi \in \hat{A}$, $e_\pi \in R_1(H_\pi)$ and $S \in X(\mathcal{H})$, we have

$$CS^*S((\pi, e_\pi)) \leq \Sigma_{j\in J} S^*S_jS_j^*S(\pi, e_\pi)) \leq DS^*S((\pi, e_\pi)),$$

so,

$$Ce_\pi \circ S_\pi^* \circ S_\pi \circ e_\pi \leq \Sigma_{j\in J} e_\pi \circ S_\pi^* \circ S_{j\pi} \circ e_\pi \circ S_{j\pi}^* \circ S_\pi \circ e_\pi \leq De_\pi \circ S_\pi^* \circ S_\pi \circ e_\pi.$$

In particular, for any nonzero element $x_\pi \in H_\pi$ we have

$$C\|S_\pi(x_\pi)\|^2 \leq \Sigma_{j\in J} \mid \langle S_\pi(x_\pi), S_{j\pi}(x_\pi)\rangle \mid^2 \leq D\|S_\pi(x_\pi)\|^2.$$

Since bounded holomorphic sections exhaust fibers, so for any $y_\pi \in K_\pi$ there exists a section $S \in X(\mathcal{H})$ such that $S_\pi(x_\pi) = y_\pi$. Thus

$$C\|y_\pi\|^2 \leq \Sigma_{j\in J} \mid \langle y_\pi, S_{j\pi}(x_\pi)\rangle \mid^2 \leq D\|y_\pi\|^2. \tag{3.1}$$

According to inequality 3.1, for all $\pi \in \hat{A}$, $0 \neq x_\pi \in H_\pi$, $0 \neq y_\pi \in K_\pi$ the following set has to be countable

$$F_{x_\pi, y_\pi} := \{j \in J : \langle y_\pi, S_{j\pi}(x_\pi)\rangle \neq 0\}.$$

In particular, if $\pi \in [W]$ then K_π is separable and so it has a countable orthonormal basis as E_π. Hence, for each $\pi \in [W]$, the following set has to be countable

$$F_{\pi,x_\pi} := \{j \in J : S_{j\pi}(x_\pi) \neq 0\} = \bigcup_{y_\pi \in E_\pi} \{j \in J : \langle y_\pi, S_{j\pi}(x_\pi)\rangle \neq 0\}.$$

Consequently, if we write $W = \{(\pi_n, e_n) : n \in \mathbb{N}\}$, then $F = \bigcup_{n\in\mathbb{N}} F_{\pi_n,x_n}$ is a countable set, where for any $n \in \mathbb{N}$, $x_n \in H_{\pi_n}$ and $e_n = x_n \otimes x_n$. Also, we use the notation $f_0 = (\pi_0, e_0)$, where $e_0 = x_0 \otimes x_0$ for some $x_0 \in H_{\pi_0}$.

For each $j \in F$, $Im(S_{j\pi_0})$ is a separable space, since H_{π_0} is separable. Then, $K_0 = \langle \bigcup_{j\in F} Im(S_{j\pi_0})\rangle$ is a separable subspace of the

non-separable Hilbert space K_{π_0}, hence there exists a unit element $y_{\pi_0} \in K_{\pi_0}$ that is orthogonal to K_0. Then for any $j \in F$, $S^*_{j\pi_0}(y_{\pi_0}) = 0$.

On the other hands, for any $j \in J \setminus F$, we have $S_{j\pi_0}(x_0) = 0$, since $(\pi_0, e_0) \in \overline{W}$ and S_j is continuous. Thus for any $j \in J$, we have $\langle y_{\pi_0} S_{j\pi_0}(x_0)\rangle = 0$. By (3.1), y_{π_0} is equal to zero, that is a contradiction. Therefore, the Hilbert A-module $X(\mathcal{H})$ admits no frames. □

4. $K(l^2) + \mathbb{C}$

In the following, we consider $A = K(H) + \mathbb{C}I_H$, where H is a separable infinite dimensional Hilbert space. Also, let $\{h_n\}_{n\in\mathbb{N}}$ be an orthonormal basis for H and $e_n = h_n \otimes h_n$, for all $n \in \mathbb{N}$.

We recall that, $\hat{A} = \{\pi_0, \pi_1\}$, where $\pi_1 = id$ and $\pi_0(T + \lambda I_H) = \lambda$, for every $T \in K(H)$ and $\lambda \in \mathbb{C}$. Thus, we can consider

$$P(A) = (\{\pi_1\} \times R_1(H)) \cup \{(\pi_0, 1)\}.$$

Note that in this case, $P(A)$ is a compact hausdorff space and also $(\pi_0, 1) \in \overline{W} \setminus W$, where $W = \{(\pi_1, e_n) : n \in \mathbb{N}\}$.

Theorem 4.1. *There exists a uniformly continuous holomorphic vector bundle of dual Hopf type over $P(A)$ satisfying the conditions of Theorem 3.1.*

Proof. Li shown that in [6, Lemma 2.1], there exists an uncountable set $\mathcal{F}$ of injective maps from $\mathbb{N}$ to $\mathbb{N}$ such that for any distinct $f, g \in \mathcal{F}$, $f(n) \neq g(n)$ for all but finitely many $n \neq N$, and $f(n) \neq g(m)$ for all $n \neq m$.

Let $K_{\pi_1} = l_2$ with the standard basis $\{z_n\}_{n\in\mathbb{N}}$ and K_{π_0} be a non-separable Hilbert space with an orthonormal basis $\{h_f\}_{f\in\mathcal{F}}$ indexed by $\mathcal{F}$. For each $f \in \mathcal{F}$ consider the isometry $u_f : H \longrightarrow l_2$, given by $u_f(h_n) = z_{f(n)}$ for all $n \in \mathbb{N}$. Also, we consider $v_f : \mathbb{C} \longrightarrow K_{\pi_0}$ by $v_f(\lambda) = \lambda h_f$.

Now, we can define $S_f : P(A) \longrightarrow (\bigcup_{e\in R_1(H)} B(H, l_2)e) \cup (B(\mathbb{C}, K_{\pi_0})1)$ by

$$S_f((\pi, e)) = \begin{cases} u_f \circ e & \pi = \pi_1 \\ v_f \circ 1 & \pi = \pi_0 \end{cases}$$

Set $V = \{\sum_{i=1}^n \lambda_i S_{f_i} : n \in \mathbb{N}, \lambda_i \in \mathbb{C}, f_i \in \mathcal{F}\}$. We claim that the function $(\pi, e) \mapsto ||S(\pi, e)||$ is continuous on $P(A)$ for every $S \in V$.

For this, we note that if $S = \sum_{i=1}^m \lambda_i S_{f_i} \in V$, then there is a finite subset J of $\mathbb{N}$ such that $f_i(n) \neq f_j(n)$, for all $n \in J^c$ and $i \neq j$. Hence,

if $e = x \otimes x$, for some unite element $x \in H$, then we have

$$\begin{aligned}||S(\pi_1, e)||^2 &= ||\sum_{i=1}^{m} \lambda_i u_{f_i}(x)||^2 = ||\sum_{i=1}^{m} \lambda_i (\sum_{n=1}^{\infty} \langle x, h_n \rangle z_{f_i(n)})||^2 \\ &= ||\sum_{i=1}^{m} \sum_{n \in J} \lambda_i \langle x, h_n \rangle z_{f_i(n)}||^2 + \sum_{i=1}^{m} ||\sum_{n \in J^c} \lambda_i \langle x, h_n \rangle z_{f_i(n)}||^2 \\ &= ||\sum_{i=1}^{m} \sum_{n \in J} \lambda_i \langle x, h_n \rangle z_{f_i(n)}||^2 + \sum_{i=1}^{m} |\lambda_i|^2 (1 - ||\sum_{n \in J} \langle x, h_n \rangle z_{f_i(n)}||^2).\end{aligned}$$

Now, if a net $\{(\pi_1, e_\alpha)\}_{\alpha \in I}$ is convergent to (π_1, e) (or $(\pi_0, 1)$) and for every $\alpha \in I$, $e_\alpha = x_\alpha \otimes x_\alpha$ for some unite element $x_\alpha \in H$, then $|\langle x_\alpha, y \rangle| \to |\langle x, y \rangle|$ (or $|\langle x_\alpha, y \rangle| \to 0$), for all $y \in H$. Consequently, for every $f \in S$ and $y_1, \cdots, y_N \in H$, we have

$$||\sum_{n=1}^{N} \langle x_\alpha, y_n \rangle z_{f(n)}||(= (\sum_{n=1}^{N} |\langle x_\alpha, y_n \rangle|^2)^{\frac{1}{2}}) \to ||\sum_{n=1}^{N} \langle x, y_n \rangle z_{f(n)}||$$

$$(\text{or } ||\sum_{n=1}^{N} \langle x_\alpha, y_n \rangle z_{f(n)}|| \to 0).$$

Thus, $||S(\pi_1, e_\alpha)|| \to ||S(\pi_1, e)||$ (or $||S(\pi_1, e_\alpha)|| \to ||S(\pi_0, 1)||$). This proves the claim.

Therefore, V is a linear space of bounded holomorphic sections with uniformly continuous norm and it exhausts each fibre. Now, as mentioned in [4], by Zorn's lemma, we can extend it to a linear space $X(\mathcal{H})$ of the bounded holomorphic sections with uniformly continuous norm, maximal with this property, and exhausting each fibre. Clearly, $X(\mathcal{H})$ satisfies the conditions of Theorem 3.1. □

The following result can be obtained from Theorems 3.1 and 4.2.

Corollary 4.2. *If $A = K(l^2) + \mathbb{C}I_{l^2}$, then there exists a Hilbert A-module that possess no frames.*

References

[1] M. Amini, M. B. Asadi, G. A. Elliott, *Frames in Hilbert C*-modules and Morita equivalent C*-algebras*, Glassgow Math., to appear.

[2] L. G. Brown, *Complements to various Stone-Weierstrass theorems for C*-algebras and a theorem of Shultz*, Comm. Math. Phys. 143 (1992), 405- 413.

[3] J. Dixmier, C*-algebras, North-Holland Publishing Company, 1977.

[4] G. A. Elliott, K. Kawamura, *A Hilbert bundle characterization of Hilbert C*-modules*, Trans. Amer. Math. Soc. 360 (2008), 4841- 4862.

[5] M. Frank, D. R. Larson, *Frames in Hilbert C*-modules and C*-algebras*, J. Operator Theory 48 (2000), 273- 314.

[6] H. Li, *A Hilbert C^*-module admitting no frames*, Bull. London Math. Soc. 42 (2010), 388- 394.
[7] R. G. Swan, *Vector bundles and projective modules*, Trans. Amer. Math. Soc. 105 (1962), 264-277.

School of Mathematics, Statistics and Computer Science, College of Science, University of Tehran, Tehran, Iran, and School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran, Iran
E-mail address: mb.asadi@khayam.ut.ac.ir

School of Mathematics, Statistics and Computer Science, Collage of Science, University of Tehran, Tehran, Iran
E-mail address: z.hasanpour@ut.ac.ir